

Abstract

Results of previous papers are used to obtain a complete classification of finite local flat commutative group schemes over mixed characteristic complete discrete valuation rings. We classify group schemes in terms of their Cartier modules. We also prove the equivalence of different definitions of the tangent space and the dimension for these group schemes. In particular we prove that the minimal dimension of a formal group law that contains a given local group scheme S as a closed subgroup is equal to the minimal number of generators for the coordinate ring of S . As an application the following reduction criteria for Abelian varieties are proved.

Let K be a mixed characteristic local field, let its residue field have characteristic p , L be a finite extension of K , let $\mathfrak{O}_K \subset \mathfrak{O}_L$ be their rings of integers. Let e be the absolute ramification index of L , $s = [\log_p(pe/(p-1))]$, e_0 be the ramification index of L/K , $l = 2s + v_p(e_0) + 1$.

For a finite flat commutative $\mathfrak{O}_L$ -group scheme H we denote the $\mathfrak{O}_L$ -dual of the module J/J^2 by TH . Here J is the augmentation ideal of the coordinate ring of H .

Let V be an m -dimensional Abelian variety over K . Suppose that V has semistable reduction over L .

Theorem (A). *V has semistable reduction over K if and only if for some group scheme H over $\mathfrak{O}_K$ there exist embeddings of H_K into $\text{Ker}[p^l]_{V,K}$, and of $(\mathfrak{O}_L/p^l\mathfrak{O}_L)^m$ into $TH_{\mathfrak{O}_L}$.*

This criterion has a very nice-looking version in the ordinary reduction case.

Theorem (B). *V has ordinary reduction over K if and only if for some $H_K \subset \text{Ker}[p^l]_{V,K}$ and M unramified over K we have $H_M \cong (\mu_{p^l,M})^m$. Here μ denotes the group scheme of roots of unity.*

Keywords: finite group scheme, Cartier module, tangent space, formal group, Abelian variety, semistable reduction, local field.

MSC 2000: 14L15, 14L05, 14G20, 11G10, 11S31.

Finite flat commutative group schemes over complete discrete valuation rings II: classification, tangent spaces, and semistable reduction of Abelian varieties

M.V. Bondarko *

December 2004

Introduction

In the paper [3] the Cartier modules of finite flat local commutative group schemes over the rings of integers of complete discrete valuation fields (defined by Oort) were used to obtain results on the generic fibre of (finite flat commutative) group schemes. As an application a certain finite (see below) p -adic good reduction criterion for Abelian varieties was proved.

In this paper we use results of the previous work and obtain a complete classification of finite local flat commutative group schemes over mixed characteristic complete discrete valuation rings in terms of their Cartier modules. We give an explicit description of the image of the Oort functor. The classification obtained has the following advantages when compared with the classification of Breuil (see [5]).

1. The modules that appear as $C(S)$ are explicitly described for arbitrary local group schemes. In [5] an explicit classification is given only for S satisfying certain extra flatness conditions; the modules for arbitrary S are obtained by extensions. We note that the subclass of group schemes that was classified in [5] explicitly also can be easily described (i.e. distinguished from other local group schemes) in terms of their Cartier modules.

*Supported by Russian Fundamental Science Foundation, grant No. 04-01-00082a. The author is deeply grateful to prof. Ju. Tschinkel and the Mathematical Department of the University of Göttingen for providing excellent working conditions.

2. The Cartier module language is Galois-stable; this is crucial for descent questions.

3. We don't restrict ourselves to the perfect residue field case.

On the other hand, the advantage of [5] is that p -group schemes there were not assumed to be local.

Our classification will probably be generalized to not necessarily local p -group schemes in a succeeding paper.

We also prove the equivalence of different definitions of the tangent spaces and the dimension of these group schemes. As an application certain finite p -adic criteria for semistable and ordinary reduction of Abelian varieties are proved. We call these criteria finite because in contrast to Grothendieck's criteria (see [7]) it is sufficient to check certain conditions on some finite p -torsion subgroup of V (instead of the whole p -torsion).

We introduce some notation.

Let K be a complete discrete valuation field of characteristic 0 with residue field of characteristic p , let L be a finite extension of K , let $\mathfrak{O}_K \subset \mathfrak{O}_L$ be their rings of integers. Let e be the absolute ramification index of L , $s = [\log_p(pe/(p-1))]$. Let $e_0 = [L : K_0]$, where K_0 is the maximal unramified subextension of L/K (for ordinary local fields e_0 is equal to $e(L/K)$), $l' = s + v_p(e_0) + 1$, $l = 2s + v_p(e_0) + 1$.

For an $\mathfrak{O}_L$ -group scheme H we denote the $\mathfrak{O}_L$ -dual of the module J/J^2 by TH . Here J is the augmentation ideal of the coordinate ring of H .

Theorem (A). *Let V be an m -dimensional Abelian variety over K . Suppose that V has semistable reduction over L .*

Then V has semistable reduction over K if and only if for a certain finite flat group scheme $H/\mathfrak{O}_K$ we have $TH_{\mathfrak{O}_L} \supset (\mathfrak{O}_L/p^l \mathfrak{O}_L)^m$ (i.e. there exists an embedding); besides there exists a monomorphism $g : H_K \rightarrow \text{Ker}[p^l]_{V,K}$.

Note that for $e < p-1$ we have $l = 1$; hence Theorem A is a generalization of Theorem 5.3 of [6] (where only the potentially good reduction case was considered).

Our technique also allows us to prove the following criteria for ordinary reduction (see subsection 5.2) easily.

Theorem (B). *Let V be an Abelian variety of dimension m over K that has good reduction over L . Then the following conditions are equivalent.*

(1) *V has good ordinary reduction over K*

(2) *For a certain multiplicative type (i.e. dual-étale) group scheme $H/\mathfrak{O}_K$ we have $TH_{\mathfrak{O}_L} \approx (\mathfrak{O}_L/p^{l'} \mathfrak{O}_L)^m$; besides there exists a monomorphism $g : H_K \rightarrow \text{Ker}[p^{l'}]_{V,K}$.*

(3) For some $H_K \subset \text{Ker}[p^{l'}]_{V,K}$ and M unramified over K we have $H_M \cong (\mu_{p^{l'},M})^m$. Here $\mu_{p^{l'},M}$ is the group scheme of $p^{l'}$ -th roots of unity.

II Let V be an Abelian variety of dimension m over K that has semistable reduction over L . Then the following conditions are equivalent.

(1) V has ordinary reduction over K .

(2) For some multiplicative type group scheme H over $\mathfrak{O}_K$ there exist embeddings of H_K into $\text{Ker}[p^l]_{V,K}$, and of $(\mathfrak{O}_L/p^l\mathfrak{O}_L)^m$ into $TH_{\mathfrak{O}_L}$.

(3) For some $H_K \subset \text{Ker}[p^l]_{V,K}$ and M unramified over K we have $H_M \cong (\mu_{p^l,M})^m$.

In the first section we remind the definition of the Cartier modules for formal groups. In this paper we will usually use the so-called invariant Cartier modules (i.e. D_F) defined in [4] and [2]. We also recall Oort's definition of the Cartier module $C(S)$ for a finite local group scheme S (see [10]); we remind that the Oort functor is fully faithful. We remind the definition of a closed submodule of a Cartier module and of a separated Cartier module. We also recall connection of closed submodules with closed subgroups (cf. [3]). We prove three technical lemmas on the Cartier modules related to the reduction of formal groups and finite group schemes over $\mathfrak{O}_L$. The lemmas are used in the proof of the main classification result (Theorem 2.1.1).

Section 2 is dedicated to the main classification theorem. We describe the image of the Oort functor; that gives us a full classification of finite flat local commutative group schemes over the rings of integers of complete discrete valuation fields. The proof that for any S the module $C(S)$ satisfies the conditions of Theorem 2.1.1 is easy. In the proof of the converse statement we present the given Cartier module as a factor of $C(F)$ for some formal group F ; next we prove that a finite height F can be chosen. The classification immediately implies that Ext^1 for local group schemes coincides with Ext^1 for their Cartier modules.

In section 3 we prove that the tangent space functor can be easily described in terms of Oort modules. This implies that the minimal dimension of a finite height formal group F that contains a given local group scheme S as a closed subgroup is equal to the minimal number of generators for the coordinate ring of S . The case $m = 1$ of this result was essentially proved in [1] (though there it was formulated in a different way). It was quite important for the program of "taming wild extensions by means of Hopf algebras". We also prove that a local group scheme of exponent p^r is truncated Barsotti-Tate iff $TH \approx (\mathfrak{O}_L/p^r\mathfrak{O}_L)^m$.

In the beginning of section 4 we remind the main generic fibre result and a certain descent result of [3]. Next we prove certain new descent statements for p -divisible groups. We prove that an Abelian variety has semistable reduction

if and only if its 'formal part' has 'good reduction'. Using these statements and results on tangent spaces we prove a certain criterion for good reduction of a potentially good reduction Abelian variety. This criterion gives a positive answer to the question of existence of a finite p -adic criterion, which B. Conrad (see [6]) attributes to N. Katz. Two more variants of this criterion were proved in [3].

In section 5 we prove Theorems A and B.

Theorem A seems to be less convenient than the corresponding finite l -adic criteria (see [13], [14]). On the other hand, being ordinary is a p -adic property. It seems to be difficult to check it l -adically; in particular, no l -adic analogue of Theorem B is known.

Note that the usage of tangent spaces makes all formulations (and certainly the proofs) much different from those in the l -adic case. In particular, we are able to apply the dimension argument to finite schemes.

One is not able to prove these criteria without using tangent spaces (even in the case $e < p - 1$).

For the convenience of the reader we note that the proofs of reduction criteria use neither the classification results of section 2 nor the technical lemmas of subsection 1.2.

The author is deeply grateful to prof. Yu. Zarhin for calling his attention to the case of semistable reduction and to prof. W. Messing for useful remarks.

Notation and conventions. We keep the notation of the introduction ($K, L, p, e, s, e_0, l, l', \mathfrak{O}_K, \mathfrak{O}_L, V, \mu_{p^r}$).

Let $\overline{L}$ denote the residue field of $\mathfrak{O}_L$, let $\mathfrak{M}$ denote the maximal ideal of $\mathfrak{O}_L$, let $\pi \in \mathfrak{M}$ be some uniformizing element of L .

We introduce $t = v_p(e) + 1$.

Let $X = (X_i) = X_1, \dots, X_m, x$ be formal variables.

F will usually denote an m -dimensional formal group law over $\mathfrak{O}_L$, $\lambda_F = (\lambda_i(X))$, $1 \leq i \leq m$ denotes the logarithm of F ; $\exp_F$ is the inverse to logarithm (with respect to composition).

In this paper a 'group scheme' will (by default) mean a finite flat commutative p -group scheme (i.e. annihilated by a power of p), $S/\mathfrak{O}_L$ means a finite flat commutative group scheme over $\mathfrak{O}_L$.

$\text{Cart} = \text{Cart}_p(\mathfrak{O}_L)$ will denote the p -Cartier ring over $\mathfrak{O}_L$ (see below); Cartier module is a module over Cart .

For a (possibly, non-commutative) ring $\mathfrak{A}$ we denote by $M_{m \times n}(\mathfrak{A})$ the module of $m \times n$ -matrices over $\mathfrak{A}$; $M_m(\mathfrak{A}) = M_{m \times m}(\mathfrak{A})$.

For finite group schemes S, T we write $S \triangleleft T$ if S is a closed subgroup scheme of T ; we write $S \subset T$ if there is a group scheme morphism $f : S \rightarrow T$

that is injective on the generic fibre.

We call the number of indecomposable summands of a finitely generated $\mathfrak{O}_L$ -module M the dimension of M .

A formal group F is called a finite height one if $[p]_F$ is an isogeny, i.e. the scheme-theoretic kernel $\text{Ker}[p]_F$ of $[p]_F : F \rightarrow F$ is a finite group scheme. This property is stable with respect to base change (see [16]). In particular, F is finite height over $\mathfrak{O}_L$ if and only if it is finite height over some $P \supset \mathfrak{O}_L$, where the residue field of P is perfect.

For a finite height formal group F and a group scheme H we write $H \triangleleft F$ if H is a closed subgroup of $\text{Ker}[p^r]_F$ for some $r > 0$ (and hence, a subgroup of the p -divisible group of F).

A group scheme or a p -divisible group (in particular, a finite height formal group) is called multiplicative type if its Cartier-dual is étale.

1 Cartier modules; some results of previous papers

In this section we introduce (a modified version of) the classical Cartier module theory for formal groups; we also recall the result of Oort and some results of the previous papers. We also prove a few technical lemmas.

1.1 The definition of the Cartier ring and the Cartier modules of formal groups

For a $\mathfrak{O}_L$ -algebra Q we denote by $\text{Cart} = \text{Cart}(Q)$ the ring that is obtained by factorising $\mathbb{Z}\langle \mathbf{f}, \langle a \rangle \rangle \langle \langle \mathbf{V} \rangle \rangle$ (here $a \in Q$, we consider non-commutative series over non-commutative polynomials) modulo the following relations:

$$\langle a \rangle \langle b \rangle = \langle ab \rangle \text{ for all } a, b \in Q; \mathbf{f}\mathbf{V} = p \quad (1)$$

$$\langle a \rangle \mathbf{V} = \mathbf{V} \langle a^p \rangle; \langle a^p \rangle \mathbf{f} = \mathbf{f} \langle a \rangle \text{ for any } a \in Q \quad (2)$$

$$\langle a \rangle + \langle b \rangle = \langle a + b \rangle + \sum_{n>0} \mathbf{V}^n \langle r_{p^n}(a, b) \rangle \mathbf{f}^n, \quad (3)$$

for all $a, b \in Q$, where r_{p^n} are the polynomials defined in [8], section 16.2. We need the property $r_{p^n}(0, x) = r_{p^n}(x, 0) = 0$.

A natural analogue of (3) is valid for $\langle a \rangle - \langle b \rangle$.

In [8] $\mathbf{V}$ was denoted by V_p , $\mathbf{f}$ was denoted by F_p .

Since Cart is $\mathbf{V}$ -complete, any finitely generated Cart -module is also $\mathbf{V}$ -complete.

If P is a Q -algebra we can define the Cart-module structure on $P[[\Delta]]$ in the following way: for $f = \sum_{i \geq 0} c_i \Delta^i \in P[[\Delta]]$, $a \in Q$ we define

$$\mathbf{V}f = f\Delta; \mathbf{f}f = \sum_{i > 0} p c_i \Delta^{i-1}; \langle a \rangle f = \sum a^{p^i} c_i \Delta^i.$$

We also remind that Cart is p -complete if Q is; if M is a $\text{Cart}(Q)$ -module, then $M/\mathbf{V}M$ has a natural structure of a Q -module defined via

$$a \cdot (x \bmod \mathbf{V}M) = \langle a \rangle x \bmod \mathbf{V}M$$

for any $x \in M$.

The main ingredients of the classical Cartier theory were the modules of curves. For a formal group F/Q and any $i > 0$ one can define the Cartier module structure of $F(x^i Q[[x]]) = \text{Ker}(F(Q[[x]]) \rightarrow F(Q[[x]]/x^i))$. The p -typical (abstract) Cartier module $C(F)$ is a certain direct $\text{Cart}(Q)$ -summand of $F(xQ[[x]])$. One can describe it explicitly in the case when Q is torsion-free.

For $h \in (\mathfrak{A}[[\Delta]])^m$, the coefficients of h are equal to $h_i = \sum_{l \geq 0} c_{il} \Delta^l$, $c_{il} \in \mathfrak{A}$, we define

$$h(x) = (h_i(x)), \quad 1 \leq i \leq m, \quad \text{where } h_i = \sum_{l > 0} c_{il} x^{p^l}. \quad (4)$$

Now let $Q = \mathfrak{O}_L$. We define $D_F = \{f \in L[[\Delta]]^m : \exp_F(f(x)) \in \mathfrak{O}_L[[x]]^m\}$. In particular, for $m = 1$ we have $\sum a_i \Delta^i \in D_F \iff \exp_F(\sum a_i x^{p^i}) \in \mathfrak{O}_L[[x]]$.

The following statements were proved in [2] and [4], also see [8].

Proposition 1.1.1. 1. $D_F \approx C(F)$ as a Cart-module.

2. $F \rightarrow D_F$ is an embedding of categories of formal groups over $\mathfrak{O}_L$ into Cart-submodules of $L[[\Delta]]^m$.

Besides, for $\dim F_i = m_i$, $i = 1, 2$ we have

$$\text{Cart}(D_{F_1}, D_{F_2}) = A \in M_{m_2 \times m_1} \mathfrak{O}_L : AD_{F_1} \subset D_{F_2}.$$

3. If $f : F_1 \rightarrow F_2$, $f \equiv AX \bmod \deg 2$, $A \in M_{m_2 \times m_1} \mathfrak{O}_L$, then the associated map $f_* : D_{F_1} \rightarrow D_{F_2}$ is the multiplication by A .

4. $D_F \equiv \mathfrak{O}_L^m \bmod \Delta$.

5. If P is the ring of integers of a complete discrete valuation field containing L then $D_{F,P} = \text{Cart}_P D_F \subset PL[[\Delta]]^m$ and $D_F = D_{F,P} \cap L[[\Delta]]^m$.

6. The map $D_F \bmod \Delta \rightarrow \{f \in xL[[x]]^m : \exp_F(f) \in \mathfrak{O}_L[[x]]^m\}/x^2 L[[x]]^m$ that is induced by (4) is an isomorphism.

The set of all possible D_F can be described by conditions $D \triangleleft L[[\Delta]]^m$ (see the definition below), $D \bmod \Delta = \mathfrak{D}_L^m$ (see [2], [3]).

One can easily see (cf. the reasoning in subsection 2.3 below) that for a large class of Cart-modules any $x_i \in M, 1 \leq i \leq m$, such that $\mathfrak{D}_L(x_i \bmod \mathbf{V}M) = M \bmod \mathbf{V}M$ generate M over Cart. Moreover, any $x \in M$ can be presented in the form $\sum_{1 \leq i \leq m} \sum_{0 \leq j} \mathbf{V}^j \langle a_{ij} \rangle x_i$.

In particular, this is valid for $M = D_F$. Hence one can choose $x_i \in D_F, x_i \equiv e_i \bmod \Delta, 1 \leq i \leq m$ ($e_i = (0, \dots, 1, \dots, 0)$ is the i -th basic vector of $\mathfrak{D}_L^m$).

1.2 Some results on Cartier modules related to the reduction

The definition of the Cartier ring is functorial. In particular, a Cart-module structure on M induces a $\text{Cart}(\overline{L})$ -modules structure on $M / \text{Cart} \langle \pi \rangle M$.

If $\overline{L}$ is perfect, the module $C(F) / \text{Cart} \langle \pi \rangle C(F)$ can be identified with the usual (covariant) Dieudonne module of the reduction of F .

We denote by F_π the formal group law $F_\pi(X, Y) = \pi^{-1}(\pi X, \pi Y)$. Its coefficients obviously belong to $\mathfrak{D}_L$.

In the proof of the main classification theorem we will need the following technical results.

Lemma 1.2.1. 1. $\text{Cart} \langle \pi \rangle D_F = \pi D_{F_\pi}$.

2. $p^s \pi D_{F_\pi} \subset \mathfrak{D}_L[[\Delta]]^m$.

Proof. 1. We can assume that F is p -typical (see [8]), i.e. the logarithm $\lambda = (\lambda_i)$ of F satisfies $\lambda_i = \sum_{1 \leq j \leq m, l \geq 0} a_{ijl} X_j^{p^l}$. Then for $b_i = (b_{ij})$; $b_{ij} = \sum a_{ijl} \Delta^l$ we have $b_i \in D_F, 1 \leq i \leq m$ (see [8]). $b_i \equiv e_i \bmod \Delta$; hence b_i generate D_F as a Cart-module. Similarly $\pi^{-1} \langle \pi \rangle b_i$ generate D_{F_π} . Since $\langle \pi \rangle b_i \in \langle \pi \rangle D_F$, we obtain $\pi D_{F_\pi} \subset \text{Cart} \langle \pi \rangle D_F$.

On the other hand, πD_{F_π} is a Cart-module. We have

$$\langle \pi \rangle \left(\sum_{1 \leq i \leq m} \sum_{0 \leq j} \mathbf{V}^j \langle a_{ij} \rangle x_i \right) = \sum_{1 \leq i \leq m} \sum_{0 \leq j} \mathbf{V}^j \langle a_{ij} \pi^{p^j-1} \rangle \langle \pi \rangle x_i \in \pi D_{F_\pi}.$$

Hence $\text{Cart} \langle \pi \rangle D_F \subset \pi D_{F_\pi}$.

2. Easy calculation; see [2]. □

For the same x_i the following statement is fulfilled.

Lemma 1.2.2. *Let $y, z \in D_F$, $z \equiv y \pmod{\pi D_{F_\pi}}$, $y = \sum_{1 \leq i \leq m} \sum_{0 \leq j} \mathbf{V}^j \langle y_{ij} \rangle x_i$. Then z can be presented in the form*

$$z = \sum_{1 \leq i \leq m} \sum_{0 \leq j} \mathbf{V}^j \langle c_{ij} \rangle x_i \text{ where } c_{ij} \equiv y_{ij} \pmod{\pi}. \quad (5)$$

Proof. We construct inductively (with respect to l) a sequence of z_l, r_l such that $z = z_l + r_l$, whence z_l satisfy the conditions of (5) with b_{ijl} staying instead of c_{ij} , and $r_l \in \mathbf{V}^l \pi D_{F_\pi}$.

We start with $z_0 = y$, $r_0 = z - y$.

Now suppose that we have $b_l, r_{l,i}$ for some $l \geq 0$. We consider the decomposition

$$r_l = \sum_{1 \leq i \leq m} \sum_{0 \leq j} \mathbf{V}^j \langle d_{ijl} \pi \rangle x_i.$$

Then according to (3) we have

$$\mathbf{V}^l \left(\sum_{1 \leq i \leq m} (\langle b_{iil} \rangle + \langle d_{iil} \pi \rangle) x_i \right) = \mathbf{V}^l \left(\sum_{1 \leq i \leq m} \langle b_{iil} + d_{iil} \pi \rangle x_i \right) + s_l$$

for certain $s_l \in \mathbf{V}^{l+1} \pi D_{F_\pi}$. Hence we may take

$$z_{l+1} = z_l + \sum_{1 \leq i \leq m} (\langle b_{iil} + d_{iil} \pi \rangle - \langle b_{iil} \rangle) x_i.$$

Since D_F is $\mathbf{V}$ -complete, passing to the limit yields the assertion. $\square$

We denote by W the twisted power series ring $\overline{L}[[\mathbf{V}]] = \sum_{i \geq 0} \mathbf{V}^i x_i$; the multiplication is defined by the rule $x \mathbf{V} = \mathbf{V} x^p$, $x \in \overline{L}$.

Let $\tilde{M}$ be a module over $\text{Cart}' = \text{Cart}(\overline{L})$ (hence also a Cart-module). Since the characteristic of $\overline{L}$ is p , we have $\mathbf{V} \mathbf{f} = p = \mathbf{f} \mathbf{V}$ in Cart' (see [8]).

Suppose that any $x \in \tilde{M}$ can be presented in the form $\sum_{1 \leq i \leq m} \sum_{0 \leq j} \mathbf{V}^j \langle y_{ij} \rangle x_i$ for $y_{ij} \in \overline{L}$. For any $z = (z_i) \in W^m$, where $z_i = \sum \mathbf{V}^j z_{ij}$, we consider

$$(z)(x) = \sum_i \sum_{j \geq 0} \mathbf{V}^j \langle z_{ij} \rangle x_i.$$

Note that $(z_1 + z_2)(x) \neq (z_1)(x) + (z_2)(x)$.

We introduce a topology on W whose basic neighbourhoods of 0 are $V^i W$. Obviously, if $s_i \rightarrow 0$ in W^m then $\sum (s_i)(x)$ converges in $\tilde{M}$.

Lemma 1.2.3. *Let X be a W -submodule of W^m ; we consider the set $Y = (X)(x) = \{(z)(x); z \in X\}$. Suppose that $\mathbf{f} x_i \in Y$ for $1 \leq i \leq m$.*

Then the following statements are fulfilled.

1. For any $y, z \in X$ there exist $u, u' \in X$ such that $(u)(x) = (y)(x) + (z)(x)$, $(u')(x) = (y)(x) - (z)(x)$.
 Besides, if $z \in \mathbf{V}^r W^m$ for $r > 0$ then we can choose $u, u' \equiv y \pmod{\mathbf{V}^r W^m}$.
2. Y is a Cart' -submodule of $\tilde{M}$.
3. For any $y \in X$, $z \in W^m \setminus X$ there exists a $u \in W^m \setminus Y$ such that $(u)(x) = (z)(x) - (y)(x)$.

Proof. 1. The proof is similar to the proof of the previous Lemma.

First we construct u .

We denote $(y)(x) + (z)(x)$ by a .

We construct inductively (with respect to l) a sequence of $b_l, r_{li} \in X$, $l, i \geq 0$, such that $\lim_i r_{li} = 0$ for any $i \geq 0$, $(b_l)(x) + \sum_i (r_{li})(x) = a$, and $r_{li} \in \mathbf{V}^l X$.

We start with $b_0 = y$, $r_{0,0} = z$, $r_{0,i} = 0$ for $i > 0$.

Now suppose that we have b_l, r_{li} for some $l \geq 0$.

For $a \in \bar{L}$, $c \in W^m$ we have $\langle a \rangle(c)(x) = (ac)(x)$. Hence

$$\langle a \rangle Y \subset Y \text{ for any } a \in \bar{L} \quad (6)$$

Now arguing similarly to the proof of Lemma 1.2.2, we obtain that we can take $b_{l+1} = b_l + \sum_{i \geq 0} r_{li}$. Indeed, according to (3)

$$(b_{l+1})(x) - (b_l)(x) - \sum_{i \geq 0} (r_{li})(x)$$

can be expanded as a sum of $(w_{l+1,i})(x)$ for some $w_{l+1,i} \in \mathbf{V}^{j_{li}} \langle \bar{L} \rangle \mathbf{f} W^m \subset \mathbf{V}^{j_{li}} X$. Here each j_{li} is greater than l , $\lim_i j_{li} = \infty$. Hence we can take $r_{l+1,i} = w_{l+1,i}$.

Lastly we take $u = \lim_l b_l$.

Obviously, for $z \in \mathbf{V}^r W^m$ this construction gives us a u that is congruent to y modulo $\mathbf{V}^r W^m$.

The construction of u' is obtained from those for u by replacing some $+$ signs by $-$.

2. We have $\mathbf{V}Y \subset Y$. Part 1 states that $Y \pm Y \subset Y$. We also have (6). It remains to check that $\mathbf{f}Y \subset Y$.

Let $d = \sum \mathbf{V}^l d_l \in W^m$; $d_l \in \bar{L}^m$. Then we have

$$\mathbf{f}(d)(x) = \sum_{l \geq 0} \mathbf{V}^l (\mathbf{f}d_l)(x).$$

Besides, $\mathbf{f}(d_l)(x) \in Y$. According to the assertion of part 1, there exists a sequence $u_i \in X$, $i \geq 0$ such that $(u_i)(x) = \sum_{0 \leq j \leq i} \mathbf{V}^j (\mathbf{f}d_j)(x)$, and $u_{i+1} \equiv u_i \pmod{\mathbf{V}^{i+1}}$. Hence there exists a limit $u = \lim u_i$; we have $(u)(x) = \mathbf{f}((d)(x))$.

Thus $\mathbf{f}Y \subset \mathbf{f}\tilde{M} \subset Y$.

3. The proof is very similar to those of of part 1. We construct inductively (with respect to l) $b_l, r_{li} \in X$, $l, i \geq 0$, such that $(b_l)(x) + \sum_i (r_{li})(x) = (z)(x) - (y)(x)$; $\lim_i r_{li} = 0$, and $r_{li} \in \mathbf{V}^l X$.

In order to pass from $b_l \notin X$ to $b_{l+1} \notin X$ we use the fact that $(W^m \setminus X) \pm X = W^m \setminus X$. $\square$

1.3 Oort modules of group schemes

Let S be a local group scheme over $\mathfrak{D}_L$; let $0 \rightarrow S \rightarrow F \rightarrow G \rightarrow 0$ be its resolution by means of finite height formal groups. We define $D(F) = \text{Coker}(C(F) \rightarrow C(G))$.

In the paper [10] the following result was proved.

Proposition 1.3.1. *$S \rightarrow C(S)$ is a well defined functor on the category of local (finite flat commutative) group schemes over $\mathfrak{D}_L$; it defines an embedding of this category into the category of Cart-modules.*

We call $C(S)$ the Oort module of S .

1.4 Closed submodules and closed subschemes

The following definition played a crucial role in [3].

Definition 1.4.1. 1. For Cart-modules $M \subset N$ we write $M \triangleleft N$, if for any $x \in M$, $\mathbf{V}x \in N$ we have $x \in M$. We call M a **closed** submodule of N .

2. Cart-module N is called **separated** if $\{0\} \triangleleft N$, i.e. N has no $\mathbf{V}$ -torsion.

For Cartier modules (i.e. modules over Cart) one can easily prove the following facts (cf. [3]).

Proposition 1.4.2. *I1. If $M_i \triangleleft M$ for $i \in I$, then $\bigcap_{i \in I} M_i \triangleleft M$.*

2. If $M_2 \triangleleft M_1$, $M_1 \triangleleft M$, then $M_2 \triangleleft M$.

3. If $M_1 \triangleleft M$, $M_2 \subset M_1$, then $M_1/M_2 \triangleleft M/M_2$.

II If $f : N \rightarrow O$ is a Cart-morphism, $M \triangleleft O$, then $f^{-1}(M) \triangleleft N$.

For a Cartier module N , $O \subset N$, we denote by $\text{Cl}_N(O)$ the smallest closed Cartier-submodule of N that contains O . In particular, $\text{Cl}_N(O) \supset \text{Cart } O$.

Obviously, if $M \subset N$ are Cart-modules, then $\text{Cl}_N(M) = M \iff \text{Cl}_{N/M}\{0\} = N/M$.

Now we remind the connection of closed modules and closed subgroup schemes.

Theorem 1.4.3. *I For a local group scheme S the module $C(S)$ is separated.*

II1. Closed submodules of $C(S)$ are in one-to-one correspondence with closed subgroup schemes of S .

2. If $M = C(S)$, $N \triangleleft M = C(H)$, where $H \triangleleft S$, then $M/N \approx C(S/H)$.

3. Conversely, exact sequence (as fppf-sheaves, i.e. the inclusion is a closed embedding) of local schemes induce exact sequences of Oort modules.

III If $f : S \rightarrow T$ is a local scheme morphism, then $\text{Ker } f_* = C(\text{Ker } f)$, where f_* is the induced Oort module homomorphism; we consider the kernel in the category of flat group schemes.

IV Let M be a closed finite index submodule of $C(F)$ where F is a finite height formal group. Then M is isomorphic to $C(G)$, where G is a formal group that is isogenous to F .

V If $M \triangleleft L[[\Delta]]^m$, $D_F \subset M$, and $M \bmod \Delta = \mathfrak{D}_L^m$, then $M = D_F$.

Parts I-III were proved in [3], IV and V were proved in [2] (in slightly different formulations).

2 The main classification theorem; extensions of group schemes

This section is dedicated to classification of group schemes in terms of their Oort modules. We describe the image of the Cartier-Oort functor, thus giving a complete classification.

2.1 Formulation

Theorem 2.1.1. *I A Cart-module M is isomorphic to $C(S)$ for S being a finite flat commutative local group scheme over $\mathfrak{D}_L$ if and only if M satisfies the following conditions.*

1. $M/\mathbf{V}M$ is a finite length $\mathfrak{D}_L$ -module.

2. M is separated.

3. $\cap_{i \geq 0} \mathbf{V}^i M = \{0\}$.

4. $M = \text{Cl}_M(\langle \pi \rangle M)$.

II The minimal dimension of a finite height formal group F such that S can be embedded into F is equal to $\dim_{\mathfrak{D}_L}(C(S)/\mathbf{V}C(S))$.

III $M = C(\text{Ker}[p^r]_F)$ for an m -dimensional formal group F if and only if in addition to the conditions of I, $p^r M = 0$ and $M/\mathbf{V}M \approx (\mathfrak{D}_L/p^r \mathfrak{D}_L)^m$.

We will denote $\dim_{\mathfrak{D}_L}(M/\mathbf{V}M)$ by $\dim M$.

2.2 The proof of necessity in Theorem 2.1.1

Let $M = C(S)$.

Suppose that $S = \text{Ker } f$, where $f : F \rightarrow G$ is an isogeny of finite height formal group laws of dimension m . Then for $M = \text{Coker } f_*$ we have $M/\mathbf{V}M = D_G/(\Delta D_G + AD_F)$ where $f \equiv AX \pmod{\deg 2}$. Since $D_F \triangleleft L[[\Delta]]^m$, we have

$$\begin{aligned} M/\mathbf{V}M &= D_G/(\Delta D_G + AD_F) \\ &= (D_G \pmod{\Delta})/(AD_F \pmod{\Delta}) \approx \mathfrak{D}_L^m/A\mathfrak{D}_L^m. \end{aligned} \quad (7)$$

We obtain that M satisfies I1. We also see that the minimal dimension of a finite height formal group F such that S can be embedded into F is not less than $\dim(C(S)/\mathbf{V}C(S))$. Besides, if $S = \text{Ker}[p^r]_F$, then $C(S)/\mathbf{V}C(S) \approx (\mathfrak{D}_L/p^r\mathfrak{D}_L)^m$.

I2 for M is part I of Theorem 1.4.3.

Since $f_*D_G \triangleleft D_F$ we obtain

$$C(S)/\mathbf{V}^r C(S) \cong D_G/(\mathbf{V}^r D_G + f_*D_F) \cong (D_G/\mathbf{V}^r D_G)/(f_*D_F/\mathbf{V}^r f_*D_F).$$

Hence

$$\varprojlim M/\mathbf{V}^r M \cong \varprojlim (D_G/\mathbf{V}^r D_G)/(f_*D_F/\mathbf{V}^r f_*D_F) = D_G/f_*D_F \cong M.$$

We immediately obtain I3 for M .

Now we prove I4. Suppose that $N = \text{Cl}_M(\langle \pi \rangle M) \neq M$. Then N corresponds to some $T \triangleleft S$. We obtain that the module $U = \{x \in D_G : x \pmod{f_*D_F} \in N\} \supset \pi D_{G_\pi}$. Since $p^r S = 0$ for some $r > 0$, we have $p^r M = 0$; hence U is a finite index submodule of D_G . Therefore U corresponds to a formal group that is isogenous to G (see Theorem 1.4.3, part IV). Since these conditions are formulated in terms of formal groups and their Cartier modules, they are not affected by complete discrete valuation fields extensions (see Proposition 1.1.1, part 5). Hence we may assume that $\overline{L}$ is perfect. We obtain that $M/\langle \pi \rangle M$ is the (covariant) Dieudonne module of the reduction of S . Since S is local, its reduction is local also. It is well known that in this case $M/\langle \pi \rangle M$ is $\mathbf{V}$ -torsion, which contradicts the existence of T .

2.3 The proof of sufficiency in Theorem 2.1.1: construction of a certain formal group

Suppose that M satisfies conditions of Theorem 2.1.1 and $\dim M = m$. We first prove that there exists a formal group F , $\dim F = m$, such that $M \approx$

D_F/C , where $C \triangleleft D_F$. Next we verify that one can choose F being a finite height formal group.

We choose representatives $l_1, \dots, l_m \in M$ for (some) $\mathfrak{D}_L$ -generators of $M/\mathbf{V}M$. We check that any $x \in M$ can be expressed as a sum

$$\sum_{1 \leq i \leq l} \sum_{0 \leq j} \mathbf{V}^j \langle a_{ij} \rangle l_i. \quad (8)$$

Indeed, for any $x \in M$ there exist $a_i \in \mathfrak{D}_L$ such that $x = \sum_{0 \leq i} \langle a_i \rangle l_i + Vx_1$ for some $x_1 \in M$. Repeating this procedure r times we obtain that

$$x = \sum_{1 \leq i \leq l} \sum_{0 \leq j < r} \mathbf{V}^j \langle a_{ij} \rangle l_i + V^r x_r$$

for some $x_r \in M$. Since $\sum \mathbf{V}^j \langle a_{ij} \rangle$ converges in Cart for any $a_{ij} \in \mathfrak{D}_L$, we obtain that there exists an $y \in M$ that can be presented in the form (8) and $x - y \in V^r M$ for any $r > 0$. Now condition I3 for M implies that $x = y$.

Next we present $\mathbf{f}l_i$ in the form (8) for all $1 \leq i \leq m$. We obtain

$$\mathbf{f}l_i = \sum_{1 \leq k \leq l} \sum_{0 \leq j} \mathbf{V}^j \langle a_{ijk} \rangle l_k. \quad (9)$$

According to section 27.7 of [8], there exists an m -dimensional formal group law F such that

$$C(F) \approx (\bigoplus \text{Cart } e_i) / \sum \text{Cart}(\mathbf{f}e_i - \sum_{1 \leq k \leq l} \sum_{0 \leq j} \mathbf{V}^j \langle a_{ijk} \rangle e_k)$$

(here e_i denote certain Cart-generating elements). We obtain that there exists a Cart-homomorphism h of $C(F)$ to M that maps e_i to l_i . Since l_i generate M over Cart, h is onto.

Since M is separated, $\text{Ker } h \triangleleft C(F)$.

2.4 Choosing a finite height formal group

Now we prove that we can choose F being a finite height group. Since the height is determined by the reduction of F , it is sufficient to consider the residues of a_{ijl} , which we denote by $\overline{a_{ijk}}$. Again we denote $M/\text{Cart}\langle\pi\rangle M$ by $\tilde{M}$. According to Lemma 1.2.2, the calculation of $\overline{a_{ijk}}$ can be done in $\tilde{M}$. We denote the images of l_i in $\tilde{M}$ by $\tilde{l}_i$.

We embed $\tilde{L}$ into a perfect field U . It is well known that F has finite height if and only if the matrix $B = b_{ik}$, where $b_{ki} = \sum_{j \geq 0} \overline{a_{ijk}} \mathbf{V}^j$ is non-degenerate over the skew-field $W' = U((V))' = \sum_{i \gg -\infty} \mathbf{V}^i c_i$. Here $W' \supset W$

is the skew-field of twisted Laurent series over U , the multiplication is defined by the rule $u^p \mathbf{V} = \mathbf{V}u$, $u \in U$.

As in subsection 1.2, for any $z = (z_i) \in W^m$ we define $(z)(\bar{l}) = \sum z_i \bar{l}_i \in \tilde{M}$.

Let $v \leq m$ be the maximal possible rank of B for all possible choices of the expansion (9).

Suppose that $v < m$. We fix the numbers $i_1, \dots, i_v$ and the corresponding columns b_{i_l} of B such that b_{i_l} are W' -independent. We obtain that all other columns belong to the W' -span of b_{i_l} for any possible choice of their coefficients. We denote $(\sum W' b_{i_l}) \cap W^m$ by X ; denote $(X)(\bar{l}_i) \subset \tilde{M}$ by Y . Since $W'(W^m) \neq \sum W' b_{i_l}$, we have $X \neq W^m$. We have $\mathbf{V}X \subset X$; $\mathbf{V}(W^m \setminus X) \subset W^m \setminus X$. Since (for some choice of F) all columns of B belong to X , we have $\mathbf{f}\bar{l}_i \in Y$ for $1 \leq i \leq m$. Hence we can apply Lemma 1.2.3 with $\bar{l}_i$ staying for x_i .

We choose a column $b = b_h$ for $h \notin \{i_l\}$. We have $b \in X$. We prove that we can modify F so that the new b will not belong to X .

We choose certain $z \in W^m \setminus X$; the choice depends on whether $Y = \tilde{M}$.

In the case $Y = \tilde{M}$ we choose any $z \in W^m \setminus X$.

Suppose that $Y \neq \tilde{M}$. According to part 3 of Lemma 1.2.3, Y is a Cart' -module. Since the closure of $\{0\}$ in $\tilde{M}$ is equal to $\tilde{M}$, we obtain that $Y \not\subset \tilde{M}$. Hence there exists $y \in \tilde{M} \setminus Y$ such that $\mathbf{V}y \in \tilde{M}$. We expand $y = (z)(\bar{l}_i)$, $z = (z_i)$, $z_i \in W$. We have $z \notin X$.

In both cases we obtain that there exists a $c \in X$ such that $(c)(\bar{l}) = \mathbf{V}(z)(\bar{l})$. According to part 3 of Lemma 1.2.3 there exists a $g \in W^m \setminus X$ such that $(g)(\bar{l}) = (c)(\bar{l}) - \mathbf{V}(z)(\bar{l}) = 0$. According to part 1 of Lemma 1.2.3, there exists a $d \in X$ such that $(d)(\bar{l}) = -(b)(\bar{l})$. Applying part 3 of Lemma 1.2.3 for $z = g$, $y = d$, we obtain that b can be replaced by $b' \notin X$.

Hence the case $v < m$ is impossible. Therefore $v = m$. Thus we can choose F so that B is invertible over W' , hence F is a finite height formal group.

Remark 2.4.1. 1. In the perfect residue field case the reasoning could be simplified by using the fact that $\mathbf{V}^u M \subset \text{Cart}(\pi)M$ for some $u > 0$. This is no longer true in the imperfect residue field case.

2. Another way of proof is possible. Suppose that for a map $f : F \rightarrow G$ of (not necessarily finite height) formal groups we have $C(G)/f_*C(F) \approx M$ for M satisfying the conditions I1 – I4. Then one can note that M can be obtained from $\text{Cart}(\pi)M$ by means of successive adjoining of roots of equations of the type $\mathbf{V}y = x$. Hence $C(\bar{G})$ can be obtained from the Cartier module for the reduction of $f_*C(F)$ by successive adjoining the roots of the equation of the type $\mathbf{V}(x_i) = (y_i)$, $1 \leq i \leq m$. Next the Oort type

reasoning could be applied to prove that $\text{Ker } f$ is a finite flat group scheme and $M = C(\text{Ker } f)$.

2.5 The end of the proof of parts I and II

We take any m -dimensional finite height F such that M can be presented as a factor of D_F (as a Cart -module).

We prove that the exponent of M is finite, i.e. for some $u > 0$ we have $p^u M = 0$. This is equivalent to $p^u D_F \subset \text{Ker } h$, where $h : D_F \rightarrow M$ is the map constructed in subsection 2.3.

First we check that for some $r > 0$ we have $p^r \tilde{M} = 0$. We recall that

$$\tilde{M} = \sum_{1 \leq i \leq m} \text{Cart}' \bar{l}_i = \text{Cl}_{\tilde{M}} \{0\}.$$

Hence there exists a sequence of Cart' -modules $\{0\} = M_0 \subset M_1 \subset M_2 \subset \dots \subset M_r = \tilde{M}$ such that for any $0 \leq i < r$ we have $M_{i+1} = M_i + \text{Cart } x_i$ for some $x_i \in M_{i+1}$ satisfying $\mathbf{V}x_i \in M_i$. We have $px_i = \mathbf{fV}x_i \in M_i$. Therefore $pM_{i+1} \subset M_i$. Thus $p^r \tilde{M} = 0$.

Next we verify that $p^g \pi D_{F_\pi} \subset \text{Ker } h$ for some $g > 0$. Since $M/\mathbf{V}M$ is a finite length $\mathfrak{D}_L$ -module, we obtain that $\text{Ker } h \bmod \Delta$ is a finite index submodule of $D_F \bmod \Delta = \mathfrak{D}_L^m$. If $x = \sum_{i \geq 0} x_i \Delta^i$, $x_i \in L^m$, then $\mathbf{fV}x - \mathbf{Vf}x = px_0$. Hence for some $d > 0$ we have $p^d \mathfrak{D}_L^m \subset \text{Ker } h$; therefore $p^d \mathfrak{D}_L[[\Delta]]^m \subset \text{Ker } h$. Using part 2 of Lemma 1.2.1 we obtain $p^{d+s} \pi D_{F_\pi} \subset \text{Ker } h$.

We have an exact sequence $0 \rightarrow \pi D_{F_\pi} / (\pi D_{F_\pi} \cap \text{Ker } h) \rightarrow M \rightarrow \tilde{M} \rightarrow 0$. Thus we can take $u = r + g$.

Since F is finite height, the module $D_F/p^u D_F$ corresponds to a finite flat local group scheme $\text{Ker}[p^u]_F$.

Since $\text{Ker } h \triangleleft D_F$, we obtain $\text{Ker } h/p^u D_F \triangleleft D_F/p^u D_F$. Hence, according to part II2 of Theorem 1.4.3, M corresponds to some group scheme S which is a factor of $\text{Ker}[p^u]_F$ by a closed subscheme.

2.6 The proof of sufficiency in Theorem 2.1.1, part III

Again we choose any m -dimensional finite height F such that M can be presented as a factor of D_F .

It is sufficient to prove that for F, h constructed above we have $p^r D_F = \text{Ker } h$. Since $p^r M = 0$, we have $p^r D_F \subset \text{Ker } h$.

Since $M/\mathbf{V}M \approx (\mathfrak{D}_L/p^r \mathfrak{D}_L)^m$, using (7) we obtain $\text{Ker } h \bmod \Delta = (p^r \mathfrak{D}_L)^m$. Hence we have $p^r D_F \subset \text{Ker } h$, $p^r D_F \bmod \Delta = \text{Ker } h \bmod \Delta$.

According to part V of Theorem 1.4.3, we have $p^{-r} \text{Ker } h = D_F$. Hence $p^r D_F = \text{Ker } h$.

2.7 Extensions of group schemes

Theorem 2.1.1 easily implies the following result for Ext^1 in the category of group schemes.

Proposition 2.7.1. *If S, T are local, then $\text{Ext}^1(S, T) = \text{Ext}_{\text{Cart}}^1(C(S), C(T))$. Here we consider extensions in the category of finite flat group schemes, whence the definition of an exact sequence is the same as always in this paper.*

Proof. Since S, T are local, any extension of T by S is also local.

We have to prove that conditions of part I of Theorem 2.1.1 are preserved by extensions of Cart-modules.

Suppose that we have an exact sequence $0 \rightarrow C(T) \rightarrow M \rightarrow C(S) \rightarrow 0$ of Cartier-modules. We check that M satisfies conditions I1 – I4.

The sequence $C(T)/\mathbf{V}C(T) \rightarrow M/\mathbf{V}M \rightarrow C(S)/\mathbf{V}C(S) \rightarrow 0$ is right-exact and we obtain I1.

Since $C(S)$ is separated, we obtain $C(T) \triangleleft M$. Hence $\{0\} \triangleleft C(T) \triangleleft M$ and we obtain I2.

Since $\cap_{i \geq 0} \mathbf{V}^i C(S) = \{0\}$, for any $x \in \cap_{i \geq 0} \mathbf{V}^i M$ we have $x \in C(T)$. Since $C(T) \triangleleft M$, we also obtain $x \in \cap_{i \geq 0} \mathbf{V}^i C(T)$. Now condition I3 for $C(T)$ implies $x = 0$. We obtain I3 for M .

Since $C(T) \subset M$, we obtain $\langle \pi \rangle C(T) \subset \langle \pi \rangle M$. Hence $\text{Cl}_M(\langle \pi \rangle M) \supset C(T) \triangleleft M$. Hence

$$\text{Cl}_M(\langle \pi \rangle M) = \{x \in M : x \bmod C(T) \in \text{Cl}_C(S)(\text{Cart} \langle \pi \rangle C(S))\} = M.$$

We obtain I4 for M .

□

3 The tangent space of a finite group scheme

3.1 Definition; expression in terms of the Oort module

We introduce a natural definition of the tangent space TS for a finite group scheme S .

Definition 3.1.1. For a finite flat group scheme S we denote by TS the $\mathfrak{D}_L$ -dual of J/J^2 (i.e. $\text{Hom}_{\mathfrak{D}_L}(J/J^2, L/\mathfrak{D}_L)$), where J is the augmentation ideal of the coordinate ring of S .

Remark 3.1.2. 1. Obviously, the tangent space is an additive functor on the category of finite group schemes.

2. The definition implies that the (suitably defined) tangent space of $S_P = S \times_{\text{Spec } \mathfrak{O}_L} \text{Spec } P$ is equal to $TS \otimes_{\mathfrak{O}_L} P$. Here P is any (unital commutative) $\mathfrak{O}_L$ -algebra.

3. It is well known that the tangent space of a group scheme is equal (i.e. naturally isomorphic) to the tangent space of its local part.

Proposition 3.1.3. 1. TS is naturally isomorphic to $C(S_0)/\mathbf{V}C(S_0)$, where S_0 is the local part of S .

2. A local group scheme homomorphism $f : S \rightarrow T$ is a closed embedding if and only if the induced map of the tangent spaces is an embedding.

3. If $0 \rightarrow H \rightarrow S \rightarrow T \rightarrow 0$ is an exact sequence of local group schemes (in the category of fppf-sheaves, i.e. $H \triangleleft S$) then the corresponding sequence of tangent spaces is also exact.

4. If $H \subset S$ then the $\mathfrak{O}_L$ -length of TH is not greater than the length of TS ; they are equal iff $H = S$.

Proof. 1. Since the local part is functorial, it is sufficient to prove the statement for $S_0 = S$. We denote the isomorphism we want to construct by i_S .

For a formal group law (i.e. for a formal Lie group) F let J_F denote the augmentation ideal of the coordinate ring of F . It is well known that J_F/J_F^2 is the cotangent space of F (at 0) i.e. we have an isomorphism

$$TF = \text{Hom}_{\mathfrak{O}_L}(J_F/J_F^2, \mathfrak{O}_L) \cong TF' = \text{Ker}(F(x\mathfrak{O}_L[[x]]) \rightarrow F(x\mathfrak{O}_L[[x]]/x^2))$$

that is canonic and functorial. Via $\log_F$ the module TF' is canonically isomorphic to $\{f \in xL[[x]]^m : \exp_F(f) \in \mathfrak{O}_L[[x]]^m\}/x^2L[[x]]^m$. Now applying part 6 of Proposition 1.1.1 we obtain $T_F \cong D_F \pmod{\Delta}$. Hence we have a functorial isomorphism $i_F : D_F/\mathbf{V}D_F \rightarrow TF$.

Let $0 \rightarrow S \rightarrow F \xrightarrow{h} G \rightarrow 0$ be a resolution of S via finite height formal groups of dimension $r > 0$. Since $h_*D_F \triangleleft L[[\Delta]]^r$, we have $h_*D_F \triangleleft D_G$. According to (7) we have

$$C(S)/\mathbf{V}C(S) = D_G/(h_*D_F + \mathbf{V}D_G) = (D_G/\mathbf{V}D_G)/h_*(D_F/\mathbf{V}D_F).$$

Besides h_* is injective on $(D_F/\mathbf{V}D_F)$. The standard schematic construction of the kernel of a formal group scheme homomorphism gives us $J \cong J_F/h^*J_G$. Moreover, $h^*J_G^2 = h^*J_G \cap J_F$. Therefore we can define i_S by means of i_G .

We note that the construction of i_S is functorial with respect to morphisms of isogenies of formal groups (i.e. with respect to commutative squares of morphisms of formal groups).

It remains to check that i_S does not depend on the choice of resolution for S and is functorial with respect to group scheme morphisms.

Let $0 \rightarrow S \rightarrow F_1 \xrightarrow{h_1} G_1 \rightarrow 0$ and $0 \rightarrow S \rightarrow F_2 \xrightarrow{h_2} G_2 \rightarrow 0$ be two different resolutions of S by means of formal groups.

It was proved in [10] that there exists a resolution $0 \rightarrow S \rightarrow F \xrightarrow{h} G \rightarrow 0$ such that for $i = 1, 2$ the following commutative diagram can be constructed:

$$\begin{array}{ccccc} S & \longrightarrow & F_i & \xrightarrow{h_i} & G_i \\ \downarrow \text{id}_S & & \downarrow & & \downarrow \\ S & \longrightarrow & F & \xrightarrow{h} & G \end{array} \quad (10)$$

the rows are exact.

Hence i_S for first two resolutions are the same as the one for the third resolution.

Now let $f : S \rightarrow T$ be a morphism of (finite flat local commutative) group schemes. Then according to [10], we can construct a commutative diagram

$$\begin{array}{ccccc} S & \longrightarrow & F_1 & \longrightarrow & G_1 \\ \downarrow f & & \downarrow & & \downarrow \\ T & \longrightarrow & F_2 & \longrightarrow & G_2 \end{array} \quad (11)$$

where the rows are resolutions of S and T . Since i_S is functorial with respect to morphisms of isogenies of formal groups, we obtain that i_S is natural.

2. If $H \triangleleft S$ then $C(H) \triangleleft C(S)$, hence we obtain that $C(H) \cap \mathbf{VC}(S) = \mathbf{VC}(H)$. Therefore the kernel of the map $C(H)/\mathbf{VC}(H) \rightarrow C(S)/\mathbf{VC}(S)$ is zero.

We prove the inverse implication.

We may decompose f as $g \circ i \circ h$, where g is a closed embedding that corresponds to the injection $\text{Im } f_L \rightarrow S_L$, i is bijective on the generic fibre, and h is epimorphic (see Proposition 4.1.1 below).

Since the induced tangent space map g_* is injective, we may assume that f_L is surjective (i.e. $g = \text{id}_S$).

If the composite $i_* \circ h_*$ of the maps of tangent spaces is injective then h_* is injective.

If we have an exact sequence $0 \rightarrow B \rightarrow H \rightarrow S \rightarrow 0$, then $\{0\} \neq TB \subset TH$; besides TB maps to 0 via the map $TH \rightarrow TS$. Hence h is an isomorphism.

It remains to consider the case $H \subset S$, $H_L = S_L$. The map $TH \rightarrow TS$ is injective iff $C(H) \cap \mathbf{VC}(S) = \mathbf{VC}(H)$. Then for $x \in C(H)$, $\mathbf{V}x \in C(S)$ we

obtain $\mathbf{V}x = \mathbf{V}y$, $y \in C(H)$. Since $C(S)$ is separated, we obtain $x = y \in C(H)$. Hence $C(H) \triangleleft C(S)$. Applying Theorem 1.4.3 we obtain $H \triangleleft S$.

3. From Theorem 1.4.3 we immediately obtain that the sequence $TH \rightarrow TS \rightarrow TT \rightarrow 0$ is right-exact. It remains to apply part 2.

4. We can assume that H and S are local.

Suppose that we have exact sequences $0 \rightarrow H_0 \rightarrow H \rightarrow H_1 \rightarrow 0$ and $0 \rightarrow S_0 \rightarrow S \rightarrow S_1 \rightarrow 0$, where $H_i \subset S_i$ for $i = 0, 1$. Then, according to the assertion of part 3, we obtain that it is sufficient to check the assumption for the pairs $H_i \subset S_i$. Applying the results of Raynaud (see Proposition 4.1.1 below) we reduce the statement to the case $H_L = S_L$, H_L is L -irreducible. Since the assumption is not affected by finite extensions of L (see Remark 3.1.2), we may also assume that the order of H is p . Then the coordinate ring of H can be presented as $\mathfrak{D}_L[x]/x^p - ax$ for some $a \in \mathfrak{D}_L$ (see the classification in [11]), the augmentation ideal is equal to (x) . Then $TH \approx \mathfrak{D}_L/a\mathfrak{D}_L$. It remains to note that $\text{Spec}(\mathfrak{D}_L[x]/x^p - ax) \subset \text{Spec} \mathfrak{D}_L[x]/x^p - bx$ implies $a \mid b$; if $b \sim a$ then the inclusion is an isomorphism. $\square$

Remark 3.1.4. One can also prove the equality $d = l\#S$ for l being the length of TS , d being the valuation of the discriminant of the coordinate ring of S . This generalizes Tate's formula for $d(\text{Ker}[p^r]_F)$ (see [15]).

3.2 The dimension of a group scheme

Now we rewrite parts II and III of Theorem 2.1.1 in terms of tangent spaces.

Theorem 3.2.1. *I For a local group scheme S the following numbers are equal.*

1. *The $\mathfrak{D}_L$ -dimension of J/J^2 .*
2. *The $\mathfrak{D}_L$ -dimension of $C(S)/\mathbf{V}C(S)$.*
3. *The minimal dimension of a finite height formal group F such that $S \triangleleft F$.*

II S is equal to $\text{Ker}[p^r]_F$ for some m -dimensional finite height formal group $F/\mathfrak{D}_L$ if and only if $p^r S = 0$ and $TS \approx (\mathfrak{D}_L/p^r \mathfrak{D}_L)^m$.

Proof. Immediate by combining Theorem 2.1.1 and Proposition 3.1.1. $\square$

It seems natural to call the number $\dim_{\mathfrak{D}_L}(J/J^2)$ the dimension of S .

Remark 3.2.2. 1. We obtain that the minimal number of generators for the coordinate ring of S is equal to the minimal possible dimension of F , $S \triangleleft F$. Obviously, the dimension of F cannot be smaller than the minimal number of generators; yet the inverse inequality is much more difficult and was not known previously.

The case $m = 1$ of this result was proved in [1] (though there it was formulated in a different way).

2. It seems very probable that the equality I1=I3 can be generalized to not necessarily local p -group schemes (and p -divisible groups). In order to do this one should modify the Cartier module theory so that it would describe not necessarily local group schemes. This would probably be done in one of the following papers.

4 Descent; a finite good reduction criterion

In the paper [3] it was proved that one can check whether an Abelian variety of potentially good reduction has good reduction over K knowing $\text{Ker}[p^{l'}]_{V,K}$. In this section we modify that criterion using the results of the previous section. We also prove that one can reduce the study of semistable reduction for an Abelian variety to the study of its 'formal part' (see subsection 4.4 below).

4.1 The generic fibre results: reminder

We often use the following fact (see [12]).

Proposition 4.1.1. *Closed subgroup schemes of $S/\mathfrak{O}_L$ are in one-to-one correspondence with closed subgroup schemes of S_L .*

In particular, for any set of $S_i \triangleleft S$ there exists a smallest (in the sense of $\subset$) group scheme H such that $S_i \triangleleft H$. H is also a closed subgroup scheme of S .

Now we remind the main result of [3]. It is a finite analogue of fullness of the generic fibre functor for p -divisible groups (proved by Tate). Besides, it implies Tate's result (see [15]) immediately.

Theorem 4.1.2. *If S, T are $\mathfrak{O}_L$ -group schemes, and $g : S_L \rightarrow T_L$ is a morphism of L -group schemes, then there exists an $h : S \rightarrow T$ over $\mathfrak{O}_L$ such that $h_L = p^s g$.*

Certainly, h is unique. One easily checks that the result is sharp, i.e. the value of s is the smallest one possible.

For $e = e_L < p - 1$ Theorem 4.1.2 implies that the generic fibre functor on the category of (finite flat commutative) group schemes is full. This is the central result of [12]; Raynaud's methods cannot be used in the case $e \geq p - 1$.

4.2 Descent lemmas: reminder

In [3] the following result was proved by means of an explicit descent reasoning for D_F .

Proposition 4.2.1. *Let F be a finite height formal group over $\mathfrak{D}_L$. Suppose that its generic fibre (as a p -divisible group) $F_L = F \times_{\text{Spec } \mathfrak{D}_L} \text{Spec } L$ is defined over K , i.e. there exists a p -divisible group Z_K over K such that*

$$Z_K \times_{\text{Spec } K} \text{Spec } L \cong F_L. \quad (12)$$

Suppose that $\text{Ker}[p^t]_Z \approx T \times_{\text{Spec } \mathfrak{D}_K} \text{Spec } K$ for some group scheme $T/\mathfrak{D}_K$; this isomorphism combined with the isomorphism (12) is the generic fibre of a certain isomorphism $T \times_{\text{Spec } \mathfrak{D}_K} \text{Spec } \mathfrak{D}_L \cong \text{Ker}[p^t]_F$. Then $Z_K \approx F'_K$ for some formal group $F'/\mathfrak{D}_K$.

4.3 Descent for p -divisible groups in terms of tangent spaces

We remind that (by definition) the dimension of a p -divisible group over $\mathfrak{D}_L$ is the dimension of its local part (as a formal group law).

Proposition 4.3.1. *Let V be a p -divisible group over K , let Y be a p -divisible group of dimension m over $\mathfrak{D}_L$. Suppose that $V \times_{\text{Spec } K} \text{Spec } L \cong Y \times_{\text{Spec } \mathfrak{D}_L} \text{Spec } L$. We denote by G the local part of Y , by J the corresponding subgroup of V .*

Then the following conditions are equivalent:

I There exists a p -divisible group Z over $\mathfrak{D}_K$ such that $J \cong Z \times_{\text{Spec } \mathfrak{D}_K} \text{Spec } K$.

II For a (finite flat commutative) group scheme $H/\mathfrak{D}_K$ we have $TH_{\mathfrak{D}_L} \approx (\mathfrak{D}_L/p'\mathfrak{D}_L)^m$; besides there exists a monomorphism $g : H_K \rightarrow \text{Ker}[p']_{V,K}$.

III We have $TH_{\mathfrak{D}_L} \supset (\mathfrak{D}_L/p'\mathfrak{D}_L)^m$ (i.e. there exists an embedding); there exists a monomorphism $g : H_K \rightarrow \text{Ker}[p']_{V,K}$.

Proof. I $\implies$ II: We can take $H = [\text{Ker } p']_{Z, \mathfrak{D}_K}$. Then, according to part II of Theorem 3.2.1, $TH_{\mathfrak{D}_L} \approx (\mathfrak{D}_L/p'\mathfrak{D}_L)^m$.

II $\implies$ III: obvious.

Now we prove III $\implies$ I. We can assume that H is local.

According to Theorem 4.1.2, there exists a morphism $h : H_{\mathfrak{D}_L} \rightarrow \text{Ker}[p^t]_Y$ whose generic fibre is equal to $p^s g$. We denote by S the kernel of h (i.e. $S \triangleleft H$; $S_L = \text{Ker } h_L$; see Proposition 4.1.1); $T = H/S$. Since h is defined over $\mathfrak{D}_K$, so are S and T .

We have an exact sequence $0 \rightarrow S_{\mathfrak{D}_L} \rightarrow H_{\mathfrak{D}_L} \rightarrow T_{\mathfrak{D}_L} \rightarrow 0$; besides $T_{\mathfrak{D}_L} \subset \text{Ker}[p^t]_G$. Since $p^s S = 0$, we obtain $p^s C(S) = 0$; hence $p^s TS = 0$. Since exact sequences of group schemes correspond to exact sequences of tangent spaces (see part 3 of Proposition 3.1.3), we obtain that $TT \supset (\mathfrak{D}_L/p^t \mathfrak{D}_L)^m$.

Hence part 4 of Proposition 3.1.3 implies that $\text{Ker}[p^t]_G = T$.

Since T and the embedding are defined over $\mathfrak{D}_K$, according to Proposition 4.2.1 we obtain that G is defined over $\mathfrak{D}_K$ also. $\square$

We also prove the following (slight) generalization of Theorem 7.5.1 of [3].

Theorem 4.3.2. *Let V be a p -divisible group over K , let Y be a p -divisible group over $\mathfrak{D}_L$. Suppose that $V \times_{\text{Spec } K} \text{Spec } L \cong Y \times_{\text{Spec } \mathfrak{D}_L} \text{Spec } L$.*

Then the following conditions are equivalent:

I. There exists a p -divisible group Z over $\mathfrak{D}_K$ such that $V \cong Z \times_{\text{Spec } \mathfrak{D}_K} \text{Spec } K$.

II. For a (finite flat commutative) group scheme $H/\mathfrak{D}_K$ we have $TH_{\mathfrak{D}_K} \approx (\mathfrak{D}_K/p^{\nu} \mathfrak{D}_K)^m$; besides there exists a isomorphism $g : H_K \rightarrow \text{Ker}[p^{\nu}]_{V,K}$.

Proof. $I \implies II$ Again we can take $H = [\text{Ker } p^{\nu}]_{Z, \mathfrak{D}_K}$.

$II \implies I$ In [3] the same statement was proved for H equal to $\text{Ker}[p^{\nu}]_U$ for some p -divisible U over $\mathfrak{D}_K$. It was noted also that it is sufficient to check the natural map $i : H/\text{Ker}[p^s]_H \rightarrow H$ is a closed embedding.

We verify that H satisfies this condition. Let T denote the local part of H . By part II of Theorem 3.2.1 T is equal to $\text{Ker}[p^{\nu}]_F$ for some formal group $F/\mathfrak{D}_K$. Hence the restriction of i to T is a closed embedding. Since the exponent of H equals p^{ν} , we have $i^{-1}(T) = T$. Since H/T is etale, i induces a closed embedding $H/T \rightarrow H/T$. Therefore i is a closed embedding itself. $\square$

4.4 Duality and semistable reduction of Abelian varieties

In order to reduce the descent question for Abelian varieties to the study of certain formal groups we will need the following statement.

Let V denote an Abelian variety that has semistable reduction over L , let V' denote the dual variety. We denote by V_p the p -torsion of V (as a p -divisible group over L), denote by V_f by the formal part of the p -torsion of V (i.e. the part corresponding to the formal group F of the Néron model of V over $\mathfrak{D}_L$). Obviously V_f corresponds to the maximal p -divisible subgroup V_p that is a generic fibre of a local scheme over $\mathfrak{D}_L$. Hence the formal part

is functorial with respect to isogenies of Abelian varieties (one could also deduce this fact from part 1 of the theorem below).

We note also that the formal part of an Abelian variety is Galois-stable over its field of definition. Hence if V is defined over K then V_f is equal to $V_{fK} \times_{\text{Spec } K} \text{Spec } L$ for a certain uniquely defined p -divisible subgroup $V_{fK} \subset V_p$ over K .

Theorem 4.4.1. *Let $V_m \subset V_f$ denote the multiplicative type part of V_p i.e. the part corresponding to the maximal multiplicative type subgroup of F . Let V'_m denote the multiplicative type part of V'_p .*

1. *The Weil pairing for V_p and V'_p induces the Cartier duality of V_p/V_f with V'_m .*

2. *Suppose also that V is defined over K . Then V has semistable reduction over K if and only if there exists a formal p -divisible group $G/\mathfrak{O}_K$ such that the generic fibre of G (as a p -divisible group) is isomorphic to V_{fK} .*

Proof. 1. We denote by V_{fin}, V'_{fin} the finite parts of V_p and V'_p respectively, by V_t, V'_t the toric parts of V_p and V'_p respectively (see the definitions in [7]). Obviously, $V'_t \subset V'_m$.

By Theorem 5.2 of [7]) V'_{fin} is the Cartier dual of V_p/V_t , V'_t is the Cartier dual of V_p/V_{fin} . Hence for D being the Cartier dual of V_p/V_f we have $V'_t \subset D \subset V'_{fin}$. Since V_{fin}/V_f is the maximal possible étale factor of V_{fin} (over $\mathfrak{O}_L$), D/V'_t is the maximal possible multiplicative type subgroup of V'_{fin}/V'_t . Hence $D = V'_m$.

2. We define V'_{fK} similarly to V_{fK} . Let $\alpha : V \rightarrow V'$ be some polarization over K . Since the formal part is functorial, α induces an isogeny $\beta : V_{fK} \rightarrow V'_{fK}$. Then $\text{Ker } \beta$ is a finite K -group scheme; hence it corresponds to some $S \triangleleft G$ (see Proposition 4.1.1). Therefore $V'_{fK} \cong Z' \times_{\text{Spec } \mathfrak{O}_K} \text{Spec } K$, where $Z' = Z/S$ is a formal p -divisible group over $\mathfrak{O}_K$.

We obtain that V'_{fK} can be defined over $\mathfrak{O}_K$. Since the multiplicative type part of an Abelian variety is Galois-stable, V'_{mK} (defined similarly to V'_{fK}) is defined over $\mathfrak{O}_K$ also. Using the assertion of part 1, we obtain that V_{fK} and V_{pK}/V_{fK} are defined over $\mathfrak{O}_K$. Hence V_{pK} is Barsotti-Tate of echelon 2 over $\mathfrak{O}_K$. By Proposition 5.13c of [7], V has semistable reduction over K . $\square$

Remark 4.4.2. In [7] V_{fin} was called the fixed part of V_p , V_m was called the toroidal part of V_p , $V_{K,fin}$ was called the effectively fixed part.

4.5 Criterion for good reduction of a potentially good reduction Abelian variety

Theorem 4.5.1. *Let V be an Abelian variety of dimension m over K that has good reduction over L . Then V has good reduction over K if and only if for a certain (finite flat commutative) group scheme $H/\mathfrak{O}_K$ we have $TH_{\mathfrak{O}_L} \supset (\mathfrak{O}_L/p'\mathfrak{O}_L)^m$ (i.e. there exists an embedding) and there exists a monomorphism $g : H_K \rightarrow \text{Ker}[p']_{V,K}$.*

Proof. If V has good reduction over K then we can take $H = [\text{Ker } p']_{Y, \mathfrak{O}_K}$, where $Y_{\mathfrak{O}_K}$ is the Néron model of V over $\mathfrak{O}_K$.

Now we prove the converse implication. By Proposition 4.3.1 the formal part of V_p is defined over $\mathfrak{O}_K$. By part 2 of Theorem 4.4.1 we conclude that V has semistable reduction over K . Hence V has good reduction over K . $\square$

In [3] we proved a version of the statement where any group scheme $H/\mathfrak{O}_K$ was allowed. We required the condition $H_K \approx \text{Ker}[p']_{V,K}$.

5 Finite criteria for semistable and ordinary reduction

5.1 Proof of Theorem A

We prove Theorem A of the introduction. We adopt the notation of Theorem 4.4.1.

If V has semistable reduction over K then we can take $H = [\text{Ker } p^l]_G$.

Now we prove the converse implication in Theorem A. We can assume that H is local.

Since the finite part of an Abelian variety is Galois-stable over its field of definition, we obtain that V_{fin} is equal to $V_{finK} \times_{\text{Spec } K} \text{Spec } L$ for a certain p -divisible group V_{finK} over K . According to Proposition 5.6 of [7], the factor V_p/V_{fin} is the generic fibre of some étale p -divisible group over $\mathfrak{O}_L$. Let H_0 denote the closed group scheme of H that corresponds to the preimage of V_{fin} in H_L (see Proposition 4.1.1). We obtain that $(H/H_0)_L$ is isomorphic to a generic fibre of a certain $\mathfrak{O}_L$ -group scheme. Since H is local, Theorem 4.1.2 implies that $p^s(H/H_0) = 0$ (in fact, this statement is very easy to prove).

We denote by H_1 the maximal closed subgroup scheme of H that is killed by p' . The same tangent space argument as in the proof of Proposition 4.3.1 shows that $TH_1 \supset (\mathfrak{O}_L/p'\mathfrak{O}_L)^m$.

We have $H_1 \triangleleft H_0$, whence $H_{0K} \triangleleft V_{finK}$. Therefore $TH_0 \supset (\mathfrak{O}_L/p'\mathfrak{O}_L)^m$. We use the same argument as in the proof of Theorem 4.5.1. By Proposition

4.3.1 the formal part of V_p is defined over $\mathfrak{O}_K$. By part 2 of Theorem 4.4.1 we conclude that V has semistable reduction over K .

5.2 Ordinary reduction: reminder

We recall that an Abelian variety (over $\mathfrak{O}_K$ or $\mathfrak{O}_L$) is called an ordinary reduction one (or just ordinary) if the formal group of its Néron module is finite height and of multiplicative type.

In particular, an ordinary variety has semistable reduction.

For example, a semistable reduction elliptic curve is either ordinary or supersingular.

One can easily see that the definition given is equivalent to the usual ones. In [9] an Abelian variety was called ordinary over $\mathfrak{O}_L$ if the connected component of 0 of the reduction of the Néron model of A over $\mathfrak{O}_L$ is an extension of a torus by an ordinary Abelian variety (over $\overline{L}$).

A good reduction Abelian variety that is ordinary is called a good ordinary reduction one.

5.3 Proof of Theorem B of the introduction

Proof. I (1) $\implies$ (2)

We can take $H = \text{Ker}[p^{l'}]_{F, \mathfrak{O}_K}$ where F is the formal group of the Néron model of V (over $\mathfrak{O}_K$).

(2) $\implies$ (1)

According to Theorem 4.5.1 if such an H exists then V has good reduction over K . The same argument as in the proof of Theorem 4.3.1 shows that $\text{Ker}[p^t]_{F, \mathfrak{O}_L}$ is multiplicative. Then F is also multiplicative. In order to see that one may pass to the duals and notice that a p -divisible group U is étale iff $\text{Ker}[p^t]_U$ is.

(2) $\implies$ (3)

Let H' denote the Cartier dual of H . Since H' is étale, H' is constant over some A unramified over $\mathfrak{O}_K$. We take M being the fraction field of A .

Since H'_A is constant, H_A is isomorphic to a $\sum_{1 \leq i \leq r} \mu_{p^{n_i}}$ for some $n_i, r > 0$. We have $\mu_{p^{n_i}, A} \cong \text{Spec } B$, where $B = A[x]/((x+1)^{p^{n_i}} - 1)$ and $J(B) = (x)$. Hence $T\mu_{p^{n_i}} \approx A/p^{n_i}A$. Since $TH_{A\mathfrak{O}_L} \approx (A\mathfrak{O}_L/p^{l'}A\mathfrak{O}_L)^m$, using the additivity of the tangent space functor we obtain that $r = m$ and $n_i = l'$ for $1 \leq i \leq r$. Thus $H_M \cong (\mu_{p^{l'}, M})^m$.

(3) $\implies$ (1)

We denote the ring of integers of M by A . We take $H_A = (\mu_{p^{l'}, A})^m$. Since $TH \approx (A/p^{l'}A)^m$, H_A satisfies the conditions of (2) over M . Hence V has

good reduction over M . Since M/K is unramified, V has good reduction over K also.

II (1) $\implies$ (2)

If V has ordinary reduction, we can take $H = \text{Ker}[p^l]_{F, \mathfrak{O}_K}$.

(2) $\implies$ (1)

We apply Theorem A. As in the proof of part I, we obtain that $\text{Ker}[p^t]_F$ is multiplicative, hence F is multiplicative. We also note that (1) and (2) are equivalent to (2) with the condition $(\mathfrak{O}_L/p^l\mathfrak{O}_L)^m \subset TH$ replaced by $TH_{\mathfrak{O}_L} \approx (\mathfrak{O}_L/p^l\mathfrak{O}_L)^m$.

(2) $\iff$ (3) The proof (for $TH_{\mathfrak{O}_L} \approx (\mathfrak{O}_L/p^l\mathfrak{O}_L)^m$) is the same as in part I.

□

Remark 5.3.1. 1. Passing to the duals, one may replace a multiplicative subgroup scheme of $\text{Ker}[p^l]_V$ by an étale factorscheme in part II(3) of Theorem B.

2. (1) $\iff$ (3) of Theorem B is quite similar to the finite l -adic semistable reduction criteria of [13] and [14].

References

- [1] Bondarko M. V., Local Leopoldt's problem for rings of integers in abelian p -extensions of complete discrete valuation fields// Doc. Math. 2000, no. 5, P. 657–693.
- [2] Bondarko M. V., Explicit classification of formal groups over complete discrete valuation fields with imperfect residue field (Russian)// Trudy St. Peterburgskogo Matematicheskogo Obsh'estva, vol. 11, P. 1–36, 2005.
- [3] Bondarko M.V., Finite flat commutative group schemes over complete discrete valuation fields I: the generic fibre functor, a finite wild criterion for good reduction of Abelian varieties// to appear in Mat. Izv. Akad. Nauk, 2006.
- [4] Bondarko M. V., Vostokov, S. V., Explicit classification of formal groups over local fields (Russian)// Tr. Mat. Inst. Steklova 241 (2003), Teor. Chisel, Algebra i Algebr. Geom., P. 43–67; translation in Proc. Steklov Inst. Math. 2003, no. 2 (241), P. 35–57.
- [5] Breuil C., Groupes p -divisibles, groupes finis et modules filtrés// Ann. of Math. 2002. vol. 152. no. 2, P. 489–549.

- [6] Conrad B., Finite group schemes over bases with low ramification// Compositio Mathematica. 1999. v. 119. P. 239–320.
- [7] Grothendieck A., Séminaire de géométrie algébrique 7 I (Exposé IX), Lecture Notes in Mathematics, vol. 288. Springer-Verlag, Berlin–Heidelberg–New York, 1972.
- [8] Hazewinkel M., Formal groups and applications. Springer-Verlag, Berlin–Heidelberg–New York, 1978.
- [9] Mazur B., Tate J., Canonical height pairings via biextensions. In: Arithmetic and geometry, Vol. I, Progr. Math., 35, Birkhäuser Boston, Boston, MA, 1983, P. 195–237,
- [10] Oort F., Dieudonné modules of finite local group schemes// Indag. Math. v. 37. 1975. P. 103–123.
- [11] Oort F., Tate J., Group schemes of prime order// Ann. Sci. École Norm. Sup. 1970. v. 3 no. 4, P. 1–21.
- [12] Raynaud M., Schemas en groupes de type $(p, \dots, p)$ // Bull. Soc. Math. France. 1974. v. 102. P. 241–280.
- [13] Silverberg A., Zarhin, Yu. G., Reduction of abelian varieties// The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), NATO Sci. Ser. C Math. Phys. Sci., 548, 2000. P. 495–513.
- [14] Silverberg A., Zarhin, Yu. G., Semistable reduction of abelian varieties over extensions of small degree// J. Pure Appl. Algebra (1998), v.132 no. 2. P. 179–193.
- [15] Tate J., p -divisible groups// Proc. Conf. Local Fields (Driebergen, 1966) P. 158–183 Springer, Berlin, 1967.
- [16] Zink Th., Cartiertheorie kommutativer formaler Gruppen. Teubner-Texte zur Mathematik 68, BSB B. G. Teubner Verlagsgesellschaft, Leipzig, 1984.